


AFIB AND HEART FAILURE TOGETHER

What Does It Mean?
And What You Can Do

ACT
against Afib

Structure and Rhythm Both Play a Role in Cardiac Output



Afib can lead to decompensation or it can be the primary stimulus for the development of HF¹

Cardiac surgeons have the golden opportunity at the time of surgery for a valve problem or coronary artery blockage to halt the progression of heart failure symptoms and restore sinus rhythm.

Atrial fibrillation (Afib) can lead to heart failure (HF) or worsen it, which carries an increased risk of stroke and death¹ – and, HF happens to be the leading cause of death in Afib patients.² However, Afib is ignored almost 80% of the time in patients already undergoing surgery^{3, 4} – even with a Class I guideline recommendation.

Evidence has proven that Afib and HF co-mingle, decreasing patients' longevity.⁵ Treating Afib in the setting of large left atria and reduced ejection fraction can improve the quality of patients' lives after surgery and help them return to a normal life expectancy curve.⁴

HF is the leading cause of death in Afib patients.²

The Imminent Epidemic of Afib and Its Concomitant Diseases

A Cause for Concern

After age 40:



1 in 4 people will develop Afib⁶



1 in 5 people will develop HF⁷



HF prevalence is expected to increase by **46%**
and costs will double to **\$70 billion** by 2030¹


Afib and HF are two conditions that are likely to dominate the next 50 years of cardiovascular care.⁸

Hiding in Plain Sight

Shared Risk Factors

Incidence of Afib and HF has been related to classic cardiovascular risk factors, generating a pro-inflammatory environment and fibrosis – a common basis of both diseases.⁹

Afib can share many of the same risk factors as HF, making it difficult to detect, and promotes HF development through multiple mechanisms.¹




Afib contributes to left ventricular dysfunction leading to HF.⁵

A person with Afib is 5x more likely to develop HF.¹⁰

The Vicious Cycle

Ignoring Afib Can Lead to HF – Or Worsen It

Previous studies and surveys have shown that common underlying symptoms **such as an enlarged left atrium and reduced ejection fraction** are perceived barriers for the treatment of Afib in cardiac surgery patients.³



What can seem like a short-term barrier to treating Afib can be a marker for long-term development or worsening of HF. Patients can still continue a negative progression following structural heart surgery.


29% of Afib patients die of HF within a decade of diagnosis.²

Benefits of Restoring Sinus Rhythm

Take advantage at the time of cardiac surgery to halt the progression of HF symptoms

Matched comparisons demonstrate that cardiac surgery patients with Afib who have surgical ablation have improved outcomes, without sacrificing perioperative safety.^{4, 11}

Patients with Afib and HF who spend a higher proportion of time in sinus rhythm suffer less severe functional impairment.⁸



Restoring sinus rhythm puts the patient back on the same life-expectancy curve as if they never had Afib.¹¹

Afib and HF Together: What Does It Mean?

It means:



Afib will accelerate progressive damage to the heart, reducing cardiac output which can cause or worsen heart failure.



Reduced quality of life, more hospitalization, and increased risk of stroke and death.



It's time to treat Afib as a **long-term risk**, taking advantage of the opportunity at the time of surgery to restore cardiac output and improving quality of life and survival for patients.

NOTES



To find out more, visit
ACTagainstAfib.com

References:

- ¹ Carlisle, M.A., Fudim, M., DeVore, A. D., & Piccini, J. P. (2019). Heart failure and atrial fibrillation, like fire and fury. *JACC: Heart Failure*, 7(6):447-56.
- ² Fauchier, L. et al. (2016). Causes of death and influencing factors in patients with atrial fibrillation. *The American Journal of Medicine*, 129(12):1278-87.
- ³ Mehaffey, J.H. et al. (2021). Barriers to atrial fibrillation ablation during mitral valve surgery. *The Journal of Thoracic and Cardiovascular Surgery*.
- ⁴ McCarthy, P.M. et al. (2021). Surgery and Catheter Ablation for Atrial Fibrillation: History, Current Practice, and Future Directions. *J Clin Med*, 11(1):210, DOI: 10.3390/jcm11010210. PMID: 35011953; PMCID: PMC8745682.
- ⁵ Gerdisch, M. (2022). The most beneficial options for patients with arrhythmia and concomitant structural heart disease: a review. *AME Surgical Journal*, 3, DOI:10.21037/asj-22-18.
- ⁶ Lloyd-Jones, D.M. et al. (2004). Lifetime risk for development of atrial fibrillation: the Framingham Heart Study. *Circulation*, 110(9):1042-46.
- ⁷ Lloyd-Jones, D.M. et al. (2002). Lifetime risk for developing congestive heart failure: the Framingham Heart Study. *Circulation*, 106:3068-72
- ⁸ Kotecha, D. & Piccini, J.P. (2015). Atrial fibrillation in heart failure: what should we do? *Eur Heart J*, 36(46):3250-7. DOI: 10.1093/eurheartj/ehv513. Epub 2015 Sep 28. PMID: 26419625; PMCID: PMC4670966.
- ⁹ Boerschel, C. S. & Schnabel, R.B. (2019). The imminent epidemic of atrial fibrillation and its concomitant diseases—myocardial infarction and heart failure—a cause for concern. *International Journal of Cardiology*, 287:162-73.
- ¹⁰ Odutayo, A. et al. (2016). Atrial fibrillation and risks of cardiovascular disease, renal disease, and deaths systematic review and meta analysis. *BMJ* 2016; 354:i4482.
- ¹¹ Mosharbash, F.N. et al. (2018). Performance of the Cox-maze IV procedure is associated with improved long-term survival in patients with atrial fibrillation undergoing cardiac surgery. *The Journal of Thoracic and Cardiovascular Surgery*, 155(1):159-170.

ATRICURE, INC.

7555 Innovation Way
Mason, Ohio 45040
Customer Service: +1 (866) 349-2342
CustomerService@AtriCure.com
www.AtriCure.com
www.ACTagainstAfib.com

PM-US-2994C-0326-G
WHY TREAT AFIB SURGICALLY?

ACT
against Afib